

Occlusion of the Right Ventricular Wall Branch of a Recessive Right Coronary Artery Resulting in Ventricular Fibrillation and Anterior ST-Segment Elevation-A Case Report

Sharma, Harish; George, Sudhakar

DOI:

[10.3389/fcvm.2020.00124](https://doi.org/10.3389/fcvm.2020.00124)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Sharma, H & George, S 2020, 'Occlusion of the Right Ventricular Wall Branch of a Recessive Right Coronary Artery Resulting in Ventricular Fibrillation and Anterior ST-Segment Elevation-A Case Report', *Frontiers in cardiovascular medicine*, vol. 7, 124. <https://doi.org/10.3389/fcvm.2020.00124>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Download date: 04. May. 2023



Occlusion of the Right Ventricular Wall Branch of a Recessive Right Coronary Artery Resulting in Ventricular Fibrillation and Anterior ST-Segment Elevation—A Case Report

Harish Sharma^{1,2*} and Sudhakar George²

¹ Institute of Cardiovascular Sciences, University of Birmingham, Birmingham, United Kingdom, ² University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom

OPEN ACCESS

Edited by:

Mohammad Reza Movahed,
University of Arizona, United States

Reviewed by:

Ismail Dogu Kilic,
Pamukkale University, Turkey
David Marti,
University of Alcalá, Spain

*Correspondence:

Harish Sharma
harish.sharma@nhs.net

Specialty section:

This article was submitted to
General Cardiovascular Medicine,
a section of the journal
Frontiers in Cardiovascular Medicine

Received: 14 April 2020

Accepted: 15 June 2020

Published: 28 July 2020

Citation:

Sharma H and George S (2020)
Occlusion of the Right Ventricular Wall
Branch of a Recessive Right Coronary
Artery Resulting in Ventricular
Fibrillation and Anterior ST-Segment
Elevation—A Case Report.
Front. Cardiovasc. Med. 7:124.
doi: 10.3389/fcvm.2020.00124

Background: Right ventricular (RV) infarction is as an extremely rare cause of isolated anterior ST-segment elevation. Occlusion of the RV branch in a recessive right coronary artery (RCA) causing isolated RV infarction and only anterior ST-elevation is extremely rare. To date, the handful of such cases reports do not describe any arrhythmia associated with this presentation. Although ventricular fibrillation (VF) has been well-documented with interruption of flow in the conus branch of the RCA, here we describe VF occurring in a patient with occlusion of the RV branch of a recessive RCA presenting with isolated anterior ST-segment elevation.

Case: A 51-year-old man presented with acute chest pain and isolated anterior ST-segment elevation on electrocardiogram (ECG). The patient developed ventricular fibrillation prior to coronary angiography requiring cardiopulmonary resuscitation. Coronary angiography revealed an unobstructed left coronary system and a recessive right coronary artery with ostial occlusion of the RV branch which was treated with a drug eluting balloon, resulting in resolution of the chest pain and ECG changes.

Conclusion: Isolated RV infarction due to RV branch occlusion can cause ECG changes mimic anterior left ventricular infarction. This presentation may be complicated by VF, even in the setting of a recessive RCA.

Keywords: RV branch occlusion, anterior ST elevation, ventricular fibrillation, myocardial infarction, right ventricular infarction

INTRODUCTION

The ST-segment of an electrocardiogram (ECG) represents the time between ventricular depolarization and repolarization. In the majority of healthy individuals, the ST-segment is isoelectric. In the context of chest pain, ST-segment elevation in the left anterior precordial leads typically reflects anteroseptal infarction due to occlusion of the left anterior descending (LAD) artery. However, there are a broad range of differentials to consider (Table 1).

Right ventricular (RV) infarction is as an extremely rare cause of isolated anterior ST-segment elevation (1–4). RV infarction is caused by obstruction of flow either in the right coronary artery (RCA) proximal to the RV branch or in the RV branch itself. Proximal occlusion of a dominant RCA typically causes RV and inferior left ventricular (LV) wall infarction and results in ST-elevation in inferior and anterior leads, although the ST elevation in the inferior leads is often masked (due to anterior ST elevation leading to reciprocal inferior ST depression). Occlusion of the RV branch in a recessive RCA causing isolated RV infarction and only anterior ST-elevation is extremely rare. To date, the handful of such cases reports do not describe any arrhythmia associated with this presentation. Although ventricular arrhythmias have been well-documented with interruption of flow in the conus branch of the RCA, here we describe ventricular fibrillation occurring in a patient with occlusion of the RV branch of a recessive RCA presenting with isolated anterior ST-segment elevation.

CASE

A 51-year-old man with a history of hypertension and cigarette smoking developed acute onset chest pain radiating to his jaw and back whilst at work. After the pain worsened over 2 h, the

patient called for an ambulance and an electrocardiogram (ECG) revealed significant anterior ST elevation (see **Figure 1**).

The patient was treated as a suspected acute ST-elevation myocardial infarction (STEMI) and taken immediately to the cardiac catheterization laboratory. Before the procedure could commence, the patient developed ventricular fibrillation (VF) requiring 20 s of cardiopulmonary resuscitation and one direct current shock before return of spontaneous circulation. Coronary angiography via the right radial artery revealed a normal left main stem (LMS) and a dominant and unobstructed left circumflex artery (LCX). The left anterior descending artery (LAD) had good antegrade flow with only mild disease in the mid-vessel (**Figures 2A–D, Video S1**). Optical coherence tomography (OCT) examination of the vessel confirmed the absence of any acute plaque rupture in the proximal and mid LAD.

Right coronary angiography revealed a recessive vessel with occlusion of the right ventricular (RV) branch at the ostium (**Figures 2E,F, 3A,C**). Flow was restored as a coronary wire was passed into the RV branch. The ostial lesion was treated with an IN.PACT Falcon 2.0 mm × 20 mm Falcon drug eluting balloon resulting in resolution of the chest pain and normalization of the ST segments (**Figures 3B,D**).

Following the procedure, the patient was transferred to the coronary care unit and was found to have a high-sensitivity troponin I of 22,880 ng/L. The following day, transthoracic echocardiography revealed normal left and right ventricular size and systolic function with no significant valvular disease. The patient experienced no further chest pain or arrhythmias and was subsequently discharged 2 days after admission.

TABLE 1 | Causes of electrocardiographic anterior ST-segment elevation.

Causes of anterior ST-segment elevation on ECG	
Coronary circulation	Anteroseptal infarction
	Spontaneous coronary artery dissection
	Prinzmetal angina
	Isolated right ventricular infarction
Inflammatory	Myocarditis
	Pericarditis
Cardiomyopathy	Takotsubo Cardiomyopathy
Metabolic	Hyperkalaemia
	Hypercalcaemia
	Hypothermia
Conduction abnormalities	Bundle branch block
	Brugada syndrome
Normal variants	Benign early repolarization

DISCUSSION

RV infarction most commonly occurs due to occlusive coronary lesions in the RCA. It is associated with inferior ST-segment elevation due to concomitant infarction of the inferior left ventricular (LV) wall. Rarely, isolated RV infarction can occur, presenting with anterior ST-segment elevation. The ECG in such patients typically demonstrates significant ST-elevation in V1–2 and smaller ST-elevation in V5–6 (5).

RV infarction in such patients can be missed and treatment delayed as time is spent assessing the LAD, which more




FIGURE 1 | ECG on admission demonstrating anterior ST-segment elevation.




FIGURE 2 | Diagnostic coronary angiography of left coronary artery system (A–D) and RCA (E,F) Views: (A,B) PA cranial; (C) PA caudal; (D) LAO Caudal; (E) LAO; (F) RAO.

frequently causes anterior ST-segment elevation when flow is interrupted. This leads to significant clinical implications as patients with RV infarction are more at risk of hypotension requiring fluid resuscitation. Isolated RV infarction causing only anterior ST-segment elevation usually occurs in the setting of a recessive RCA. In this coronary pattern, the left circumflex artery

supplies the inferior LV wall and the recessive RCA supplies only the RV free wall. Acute occlusion of the recessive RCA (or its RV branch), causes isolated RV infarction and the right ventricular electrical potentials are recorded in the left precordial leads producing anterior ST-segment elevation. The reason for this lies in the anatomical position of the RV which overlies the




FIGURE 3 | Views of the RCA before (A,C) and after (B,D) drug eluting balloon treatment. Markers indicate the RV branch which was occluded and subsequently recanalized following intervention.

LV and thus infarction of the RV is recorded particularly in the first two precordial leads of the ECG (as seen in **Figure 1**).

While this presentation is rare, the particularly novel finding of this case is the occurrence of VF. Ventricular arrhythmias have been well-documented when the blood flow to the conus (first) branch of the RCA is interrupted (e.g., iatrogenic intubation of the conus branch by a coronary catheter during angiography and injection of contrast dye). The conus branch supplies the RVOT and ischaemia in this myocardial region can generate electrical disturbances and even VF. Such ventricular arrhythmias occurring due to occlusion of the RV (2nd) branch of a recessive RCA is extremely rare. Clinicians should be mindful that a patient presenting with isolated RV branch occlusion in a recessive RCA can present with only anterior ST-segment elevation and can develop life-threatening ventricular arrhythmias.

DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article/**Supplementary Material**.

ETHICS STATEMENT

Written informed consent was obtained from the individual (who is over the age of 16) for the publication of any potentially identifiable images or data included in this article.

AUTHOR CONTRIBUTIONS

Both authors contributed to the writing and proof-reading of the manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcvm.2020.00124/full#supplementary-material>

Figure S1 | Post-procedure ECG showing residual anterior ST-segment elevation.

Video S1 | Left coronary angiogram (PA cranial view).

Video S2 | Right coronary angiogram (RAO view) before intervention.

Video S3 | Right coronary angiogram (RAO view) after intervention.

REFERENCES

1. Patel A, Shenoy MM, Rao H. Case report: isolated right coronary artery obstruction presenting as acute anterior myocardial infarction. *Clin Radiol.* (1991) 44:432–3.
2. Porter A, Herz I, Strasberg B. Isolated right ventricular infarction presenting as anterior wall myocardial infarction on electrocardiography. *Clin Cardiol.* (1997) 20:971–3.
3. Sidhu MS, Aggarwal K, Fay WP. Acute isolated right ventricular myocardial infarction masquerading as acute anterior myocardial infarction. *BMJ Case Rep.* (2014) 2014:bcr2012008087. doi: 10.1136/bcr-2012-008087
4. Gregory SA, Desai AS, Fifer MA. Isolated right ventricular infarction resulting from occlusion of a nondominant right coronary artery. *Circulation.* (2004) 110:e500–1. doi: 10.1161/01.CIR.0000147608.81302.D3
5. Geft IL, Shah PK, Rodriguez L, Hulse S, Maddahi J, Berman DS, et al. ST elevations in leads V1 to V5 may be caused by right coronary artery occlusion and acute right ventricular infarction. *Am J Cardiol.* (1984) 53:991–6.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Sharma and George. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.